

# Exploring Granger Causality for Time series via Wald Test on Estimated Models with Guaranteed Stability


Nuntanut Raksasri  
Jitkomut Songsiri

Department of Electrical Engineering, Faculty of Engineering,  
Chulalongkorn University

Nov 3, 2016

- Introduction and Objective
- Granger causality
- Statistical test for Granger causality Analysis
- Stability Conditions
- Result and Conclusion


# Introduction and Objective



By knowing the relationship of the parameters in time series data, we can explain the dynamic of the time series data.

- explain relationship between the time series data by using the Granger causality concept and autoregressive model.
- estimate stable model with Granger causality constraints.

# Flow Chart of the Paper



# Auto-Regressive (AR) Model

Time series data can be represented by an AR model,

$$y(t) = c + A_1y(t-1) + A_2y(t-2) + \dots + A_p y(t-p) + v(t) \quad (1)$$

- $y(t) = (y_1(t), y_2(t), \dots, y_n(t)) \in \mathbb{R}^n$
- $A_1, A_2, \dots, A_p \in \mathbb{R}^{n \times n}$  are AR coefficients ( $p$  is lag order of the model)
- $c \in \mathbb{R}^n$  is a constant vector
- $v(t)$  is a Gaussian noise process with variance  $\Sigma$
- the observations  $y(1), y(2), \dots, y(N)$  are available.
- $y(1), y(2), \dots, y(p)$  are deterministic values and given.

“Granger causality” is a term for a specific notion of causality in time-series analysis. The idea of Granger causality is a simple one:

$$X \xrightarrow{G\text{-causes}} Y$$

**A variable  $X$  “Granger-causes”  $Y$  if  $Y$  can be better predicted using the histories of both  $X$  and  $Y$  than it can using the history of only  $Y$ .**

- Apply the concept of Granger causality to AR model in equation (1)

the causality of the model can be written in linear equation form that is if  $y_j$  “not Granger-cause” to  $y_i$  then

$$(A_k)_{ij} = 0, \quad k = 1, 2, \dots, p$$


where  $(A_k)_{ij}$  denotes the  $(i, j)$  entry of  $A_k$ .

- Granger Causality structure can be read from the zero pattern of estimated AR coefficient matrix.

## example

Consider AR(4) (AR model when  $p=4$ ) and  $y(t) \in \mathbb{R}^3$ ,  
if  $y_2$  “not Granger-cause” to  $y_1$  then the model could be

$$\begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix} = c + \begin{bmatrix} X & 0 & X \\ X & X & X \\ X & X & X \end{bmatrix} \begin{bmatrix} y_1(t-1) \\ y_2(t-1) \\ y_3(t-1) \end{bmatrix} + \begin{bmatrix} X & 0 & X \\ X & X & X \\ X & X & X \end{bmatrix} \begin{bmatrix} y_1(t-2) \\ y_2(t-2) \\ y_3(t-2) \end{bmatrix} \\ + \begin{bmatrix} X & 0 & X \\ X & X & X \\ X & X & X \end{bmatrix} \begin{bmatrix} y_1(t-3) \\ y_2(t-3) \\ y_3(t-3) \end{bmatrix} + \begin{bmatrix} X & 0 & X \\ X & X & X \\ X & X & X \end{bmatrix} \begin{bmatrix} y_1(t-4) \\ y_2(t-4) \\ y_3(t-4) \end{bmatrix} + v(t)$$





# Maximum likelihood estimation

To estimate  $A$  and  $\Sigma$  by using maximum likelihood estimation, we solving the problem

$$\underset{A, \Sigma}{\text{maximize}} \quad \frac{N-p}{2} \log \det \Sigma^{-1} - \frac{1}{2} \|L(Y - AH)\|_F^2 \quad (2)$$

when  $L^T L = \Sigma^{-1}$  and the problem is equivalent to the least-squares problem

$$\underset{A}{\text{minimize}} \quad \|Y - AH\|_F^2 \quad (3)$$

where

$$Y = [y(p+1) \quad y(p+2) \quad \cdots \quad y(N)]_{n \times (N-p)},$$
$$H = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ y(p) & y(p+1) & \cdots & y(N-1) \\ y(p-1) & y(p) & \cdots & y(N-2) \\ \vdots & \vdots & & \vdots \\ y(1) & y(2) & \cdots & y(N-p) \end{bmatrix}_{(np+1) \times (N-p)}$$

# Statistical test for Granger causality Analysis

The null hypothesis for Granger causality condition will be

$$H_0 : (A_k)_{ij} = 0 \quad \text{for } k = 1, 2, \dots, p$$

and the Wald test is based on the following test statistic:

$$W_{ij} = \hat{B}_{ij}^T \left[ \widehat{\mathbf{Avar}}(\hat{\theta})_{ij} \right]^{-1} \hat{B}_{ij}$$


where  $\hat{B}_{ij} = \left( (\hat{A}_1)_{ij}, (\hat{A}_2)_{ij}, \dots, (\hat{A}_p)_{ij} \right)$ ,  $\hat{\theta}$  is the vectorization of  $\hat{A}$ , and  $\widehat{\mathbf{Avar}}(\hat{\theta})_{ij}$  is the main diagonal block of a consistent estimate of the asymptotic covariance matrix of  $\hat{\theta}$ .

# Statistical test for Granger causality Analysis

IDEA : if  $H_0$  is true,  $(\hat{A}_k)_{ij}$  should equal to 0

In Wald test,  $H_0$  is reject if  $W > C$

where  $C = F^{-1}(1 - \alpha)$  is critical value  
and  $\alpha = \text{Prob}(W > C)$  is the significance level.



If  $W_{ij} > C \longrightarrow$  Non Zero

$C = 9.4877$  when  $\alpha = 0.05$  and  $p = 3$

**W**

X	3.2486	35.0241	0.1986	11.5292
0.1279	X	153.831	35.5729	3.2287
13.4569	0.608	X	15.0093	2.1324
12.3471	21.6202	30.7289	X	2.2625
43.4160	2.3035	2.2536	1.5259	X


**Zeros Structure**

X	0	X	0	X
0	X	X	X	0
X	0	X	X	0
X	X	X	X	0
X	0	0	0	X


Under the null hypothesis that  $(A_k)_{ij} = 0$ , the Wald statistic  $W$  converges in distribution to Chi-square distribution with  $p$  degrees of freedom.

# Wald test Results

By generating  $y(t) = AH(t) + v(t)$  and the model parameter  $A_k$  are generated by choosing some element to be equal to zero.



(a)  $\alpha = 0.01$



(b)  $\alpha = 0.1$

□ is intersect between non-zeros components

○ is true component is zero but the estimated is non-zero

+ is true component is non-zero but the estimated is zero

and blank is intersect between zero components

# AR estimation with Granger causality constraints

After knowing the Granger causality pattern, we solve this optimization problem to estimate  $A$  with Granger causality condition.

$$\begin{aligned} & \underset{A}{\text{minimize}} && \|Y - AH\|_F^2 \\ & \text{subject to} && (A_k)_{ij} = 0 \end{aligned}$$

- The estimated model parameter is not guaranteed to be stable
- we need a stability condition.

# Stability Condition

we can write the AR model in discrete-time linear system

$$\begin{bmatrix} y(t) \\ y(t-1) \\ \vdots \\ y(t-p+1) \end{bmatrix} = \underbrace{\begin{bmatrix} A_1 & A_2 & \dots & A_{p-1} & A_p \\ I & 0 & \dots & 0 & 0 \\ 0 & I & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & I & 0 \end{bmatrix}}_{\mathcal{A}} \begin{bmatrix} y(t-1) \\ y(t-2) \\ \vdots \\ y(t-p+1) \\ y(t-p) \end{bmatrix}$$

The system is stable if and only if  $\max_i |\lambda_i(\mathcal{A})| < 1$ . The characteristic polynomial have  $A_k$  as a coefficient so the condition will be nonlinear in  $A$

## Previous Works of Stability condition

Researcher	Method	Difference in paper
J. Mari et al., 2000 S. L. Lacy et al., 2003	Apply Lyapunov theory to vector ARMA	Too complicate
V. Cerone et al., 2010	Jury's test to guarantee BIBO stability on SISO LTI system	MIMO linear system
L. Buesing et al., 2012	Apply Lyapunov theory on LDS model to estimate $A$ that comply with $A^T A \prec I$	structure of $A$
K. Turksoy et al., 2013	Use Gershgorin circle theory on ARMAX	similar

# Sufficient Condition for stability

## Spectral radius and Induced norm

From spectral radius  $\rho(\mathcal{A}) = \max_i |\lambda_i(\mathcal{A})|$  then the system is stable if  $\rho(\mathcal{A}) < 1$  and by the inequality

$$\rho(\mathcal{A}) \leq \|\mathcal{A}\|$$

if assume that  $\|\mathcal{A}\| < 1$  it will affect  $\rho(\mathcal{A}) < 1$  when  $\|\mathcal{A}\|$  is a induced norm

We choose the infinity-norm of  $\mathcal{A}$  to be the sufficient condition for stability.

$$\|\mathcal{A}\|_\infty \leq 1$$

Due to structure of  $\mathcal{A}$

- $\|\mathcal{A}\|_1 \leq 1$  and  $\|\mathcal{A}\|_2 \leq 1$  lead to meaningless which is  $A_1, \dots, A_{p-1}$  are equal to zero.
- $\|\mathcal{A}\|_F \leq 1$  is impossible



We can estimate a stable model parameter with Granger causality condition by solving this problem.


$$\underset{A}{\text{minimize}} \quad \|Y - AH\|_F^2$$

$$\text{subject to } \mathcal{A} = \begin{bmatrix} A_1 & A_2 & \dots & A_{p-1} & A_p \\ I & 0 & \dots & 0 & 0 \\ 0 & I & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & I & 0 \end{bmatrix},$$


$$\|\mathcal{A}\|_\infty \leq 1,$$

$$(A_k)_{ij} = 0, \quad (i, j) \in \{1, \dots, n\} \times \{1, \dots, n\}$$

The problem is convex in quadratic form and can be solved by many solver.



(c) no constraint



(d) with Granger causality constraints

**Figure 1:** Positions of the eigenvalues of  $\mathcal{A}$  on complex plane










Figure 2: Positions of the eigenvalues of  $\mathcal{A}$  on complex plane with Granger causality and stability constraints

# Conclusion



## Why we have to check the stability?



-  L. Buesing, J. H. Macke, and M. Sahani.  
Learning stable, regularised latent models of neural population dynamics.  
*Network: Computation in Neural Systems*, 23(1-2):24–47, 2012.
-  V. Cerone, D. Piga, and D. Regruto.  
Bounding the parameters of linear systems with stability constraints.  
In *American Control Conference (ACC), 2010*, pages 2152–2157.  
IEEE, 2010.
-  W. H. Greene.  
Econometric analysis, 2008.  
*Upper Saddle River, NJ: Prentice Hall*, 24:25–26.
-  J. Mari, P. Stoica, and T. McKelvey.  
Vector arma estimation: A reliable subspace approach.  
*Signal Processing, IEEE Transactions on*, 48(7):2092–2104, 2000.



S. L. Lacy and D. S. Bernstein.

Subspace identification with guaranteed stability using constrained optimization.

*Automatic Control, IEEE Transactions on*, 48(7):1259–1263, 2003.



J. Songsiri.

Sparse autoregressive model estimation for learning granger causality in time series.

*In Acoustics, Speech and Signal Processing (ICASSP), 2013 IEEE International Conference on*, pages 3198–3202. IEEE, 2013.



K. Turksoy, E. S. Bayrak, L. Quinn, E. Littlejohn, and A. Cinar.

Guaranteed stability of recursive multi-input-single-output time series models.

*In American Control Conference (ACC), 2013*, pages 77–82. IEEE, 2013.

Q&A



THANK YOU